

Production of resonances in a thermal model^a

W. Broniowski¹, W. Florkowski^{1,2}, and B. Hiller³

¹ The H. Niewodniczański Institute of Nuclear Physics, Polish Academy of Sciences, PL-31342 Kraków, Poland

² Institute of Physics, Świętokrzyska Academy, PL-25406 Kielce, Poland

³ Centro de Física Teórica, University of Coimbra, P-3004 516 Coimbra, Portugal

Received

Abstract. We discuss the $\pi^+\pi^-$ invariant-mass correlations and the resonance production in a thermal model with single freeze-out. The predictions are confronted with the data from the STAR Collaboration.

Keywords: heavy-ion collisions, statistical model, production of resonances, particle correlations

PACS: 25.75.-q, 24.85.+p

^aPresented at the *3rd Budapest Winter School on Heavy-Ions*, 8-11 Dec. 2003.

Hadronic resonances, seen in correlation experiments at RHIC, provide valuable information on the mechanisms of particle production in relativistic heavy-ion collisions. In this talk we present the $\pi^+\pi^-$ invariant-mass correlations obtained in the framework of the *single-freeze-out model* of Ref. [1]. More details of this analysis are contained in Ref. [2].

Here is our *single-freeze-out* thermal model in a capsule:

1. The assumption of a single freeze-out, $T_{\text{chem}} = T_{\text{kin}} \equiv T$, is made.
2. We incorporate all particles from the Particle Data Tables, with their experimental branching ratios.
3. The freeze-out hypersurface is assumed to have the shape similar to the Buda-Lund model [3] (see also [4]). It possesses a Hubble-like flow, $u^\mu = x^\mu/\tau$.
4. The model has four parameters which depend on the collision energy. The two thermal parameters, $T = 160$ MeV and $\mu_B = 26$ MeV at $\sqrt{s_{NN}} = 200$ GeV, are fixed with the help of ratios of the particle abundances. The geometric parameters (the invariant time τ and the transverse size ρ_{max}) depend on centrality. They are determined by fitting the transverse-momentum spectra [1].




Fig. 1. The $\pi^+\pi^-$ invariant-mass correlation from decays of resonances in a thermal model with a static source. (a) $T_{\text{therm}} = T_{\text{chem}} = 165$ MeV. (b) $T_{\text{therm}} = 110$ MeV. The numbers indicate the position of the ρ peak (in MeV) and the labels indicate the resonance.

In order to analyze the $\pi^+\pi^-$ invariant mass spectra, measured by the STAR collaboration [5] with the like-sign subtraction technique, we assume that all the correlated pion pairs originate from the decays of neutral resonances. We include ρ , K_S^0 , ω , η , η' , f_0/σ , and f_2 . The phase-shift formula for the volume density of resonances with spin degeneracy g has the form [6]

$$\frac{dn}{dM} = g \int \frac{d^3p}{(2\pi)^3} \frac{d\delta_{\pi\pi}(M)}{\pi dM} \left[\exp\left(\frac{\sqrt{M^2 + p^2}}{T}\right) - 1 \right]^{-1}. \quad (1)$$

It was used in Ref. [2]; the same formalism has been also applied by Pratt and Bauer [7] in a similar analysis.

We first show a very simple calculation where a static source is used. Fig. 1 shows the mid-rapidity invariant-mass spectra computed at two different values of the freeze-out temperature: $T = 165$ MeV and 110 MeV. We note a clear appearance of the included resonances, as well as a significantly steeper fall-off of the strength in Fig. 1(b), which simply reflects the lower temperature. In that sense the $\pi^+\pi^-$ invariant-mass spectra provide yet another hadronic “*thermometer*” for the temper-




Fig. 2. Single-freeze-out model vs. the data of Ref. [5] for the $\pi^+\pi^-$ invariant-mass correlations. The model calculation includes the decays of higher resonances, the flow, the kinematic cuts, and the detector efficiency.

ature at freeze-out. The collective flow affects the measured resonance yields due to the presence of kinematic cuts. We now show the full calculation in our model, which includes the flow, kinematic cuts, and decays of higher resonances. The complete results are contained in Ref. [2]. We have found that the thermal effects and the kinematic cuts are able to shift down the position of the ρ peak by about 30 MeV, leaving a few tens of MeV unexplained. For that reason, we have done the calculation with the position of the ρ peak moved down (*cf.* Ref. [8]) by 9%. In Fig. 2 we compare the model predictions to the STAR data. Our model results have been filtered by the detector efficiency correction. We note that the model does an excellent job in reproducing the gross features of the data. We also note that the full model, with feeding from higher resonances and with flow and cuts, which uses the standard single-freeze-out temperature $T = 165$ MeV, gives similar predictions to the naive model at $T = 110$ MeV. This is yet another manifestation of the effect of “cooling” induced by the resonances [9]. Table 1 shows our predictions for yields of several resonances for two cases of the ρ -meson mass: 770 MeV and 700 MeV. References to the experimental data can be found in Ref. [2].

WB and WF are grateful to the organizers of the conference for providing an excellent scientific and social atmosphere, and to Patricia Fachini for many helpful discussions, for filtering our results with the STAR detector efficiency, and for preparing Fig. 2. This research was supported in part by the Polish State Committee for Scientific Research grant 2 P03B 059 25, and by the grants of Fundação para a Ciência e a Tecnologia, POCTI/FIS/35304/2000 and PRAXIS XXI/BCC/429/94.

	$m_\rho^* = 770 \text{ MeV}$	$m_\rho^* = 700 \text{ MeV}$	Experiment
$T \text{ [MeV]}$	165.6 ± 4.5	167.6 ± 4.6	
$\mu_B \text{ [MeV]}$	28.5 ± 3.7	28.9 ± 3.8	
η/π^-	0.120 ± 0.001	0.112 ± 0.001	$0.183 \pm 0.028 \text{ (40-80\%)}$
ρ^0/π^-	0.114 ± 0.002	0.135 ± 0.001	
ω/π^-	0.108 ± 0.002	0.102 ± 0.002	
$K^*(892)/\pi^-$	0.057 ± 0.002	0.054 ± 0.002	
ϕ/π^-	0.025 ± 0.001	0.024 ± 0.001	
η'/π^-	0.0121 ± 0.0004	0.0115 ± 0.0003	
$f_0(980)/\pi^-$	0.0102 ± 0.0003	0.0097 ± 0.0003	$0.042 \pm 0.021 \text{ (40-80\%)}$
$K^*(892)/K^-$	0.33 ± 0.01	0.33 ± 0.01	$0.205 \pm 0.033 \text{ (0-10\%)}$ $0.219 \pm 0.040 \text{ (10-30\%)}$ $0.255 \pm 0.046 \text{ (30-50\%)}$ $0.269 \pm 0.047 \text{ (50-80\%)}$
$\Lambda(1520)/\Lambda$	0.061 ± 0.002	0.062 ± 0.002	$0.022 \pm 0.010 \text{ (0-7\%)}$ $0.025 \pm 0.021 \text{ (40-60\%)}$ $0.062 \pm 0.027 \text{ (60-80\%)}$
$\Sigma(1385)/\Sigma$	0.484 ± 0.004	0.485 ± 0.004	

References

1. W. Broniowski and W. Florkowski, Phys. Rev. Lett. **87** (2001) 272302; Phys. Rev. C **65** (2002) 024905; W. Broniowski, A. Baran, and W. Florkowski, Acta Phys. Pol. B **33** (2002) 4235; A. Baran, W. Broniowski, and W. Florkowski, Acta Phys. Pol. B **35** (2004) 779.
2. W. Broniowski, W. Florkowski, and B. Hiller, Phys. Rev. C **68** (2003) 034911.
3. T. Csörgő and B. Lörstad, Phys. Rev. C **54** (1996) 1390; T. Csörgő, L. P. Csernai, Y. Hama, and T. Kodama, nucl-th/0306004; T. Csörgő, Heavy Ion Phys. **15** (2002) 1; M. Csanád, T. Csörgő, and B. Lörstad, nucl-th/0310040; M. Csanád and T. Csörgő, these proceedings.
4. Y. M. Sinyukov, S. V. Akkelin, and Y. Hama, Phys. Rev. Lett. **89** (2002) 052301.
5. J. Adams et al., STAR Collaboration, nucl-ex/0307023; P. Fachini, STAR Collaboration, Acta Phys. Polon. **B35** (2004) 183.
6. R. Dashen, S. Ma, and H. J. Bernstein, Phys. Rev. **187** (1969) 345; R. F. Dashen and R. Rajaraman, Phys. Rev. D **10** (1974) 694; *ibid.* p. 708; W. Weinhold, B. L. Friman, and W. Nörenberg, Acta Phys. Pol. B **27** (1996) 3249; Phys. Lett. B **433** (1998) 236; W. Weinhold, *Thermodynamik mit Resonanzzuständen*, Dissertation, TU Darmstadt, 1998; K. G. Denisenko and S. Mrówczyński, Phys. Rev. C **35** (1987) 1932.
7. S. Pratt and W. Bauer, Phys. Rev. C **68** (2003) 064905.
8. E. V. Shuryak and G. E. Brown, Nucl. Phys. A **717** (2003) 322; P. F. Kolb

-
- and M. Prakash, Phys. Rev. C **67** (2003) 044902; R. Rapp, hep-ph/0305011.
9. W. Florkowski, W. Broniowski, and M. Michalec, Acta Phys. Pol. B **33** (2002) 761.